


# Intermediate dimensions of Bedford–McMullen carpets

Amlan Banaji<sup>1</sup>

University of St Andrews



<sup>1</sup>Based on joint work with István Kolossváry in <https://arxiv.org/abs/2111.05625>

# Intermediate dimensions

- Box dimension: cover a set with balls of the same size.
- Hausdorff dimension: sets in the cover can have different sizes:


$$\dim_{\mathrm{H}} F \leq \overline{\dim}_{\mathrm{B}} F$$

- Intermediate dimensions (Falconer, Fraser and Kempton, '20) for  $\theta \in (0, 1)$ :

$$\overline{\dim}_{\theta} F = \inf \{ s \geq 0 : \text{for all } \epsilon > 0 \text{ there exists } \delta_0 \in (0, 1] \text{ such that for all } \delta \in (0, \delta_0) \text{ there exists a cover } \{U_1, U_2, \dots\} \text{ of } F \text{ such that } \delta^{1/\theta} \leq |U_i| \leq \delta \text{ for all } i, \text{ and } \sum_i |U_i|^s \leq \epsilon \}.$$


# Bedford–McMullen carpets

Divide a square into an  $m \times n$  grid,  $2 \leq m < n$ .




# Bedford–McMullen carpets

Divide a square into an  $m \times n$  grid,  $2 \leq m < n$ .




# Bedford–McMullen carpets

Divide a square into an  $m \times n$  grid,  $2 \leq m < n$ .




# Bedford–McMullen carpets

Divide a square into an  $m \times n$  grid,  $2 \leq m < n$ .



# Bedford–McMullen carpets

Divide a square into an  $m \times n$  grid,  $2 \leq m < n$ .



# Hausdorff and box dimensions




Figure: Three different Bedford–McMullen carpets, by Jonathan Fraser

Bedford ('84) and McMullen ('84) independently calculated their Hausdorff and box dimensions.

Throughout, we assume that  $\Lambda$  has **non-uniform vertical fibres**, or equivalently that  $\dim_{\text{H}} \Lambda < \dim_{\text{B}} \Lambda$ .



# Graph of the intermediate dimensions






Figure: Here,  $\gamma := \log n / \log m$


Proof involves constructing a cover using an increasing number of scales as  $\theta \rightarrow 0$ .

# Intermediate dimensions of Bedford–McMullen carpets


- Phase transitions at negative integer powers of  $\log n / \log m$ .
- Real analytic and strictly concave between phase transitions
- Strictly increasing
- Right derivative tends to  $\infty$  as  $\theta \rightarrow 0$
- Continuous for  $\theta \in [0, 1]$  (Falconer–Fraser–Kempton, '20)




# Different possible shapes of the graph



# Different possible shapes of the graph



# Different possible shapes of the graph



# Multifractal analysis

- Let  $\nu$  be the uniform Bernoulli measure supported on a Bedford–McMullen carpet, satisfying

$$\nu(A) = \sum_{i=1}^N \frac{1}{N} \nu(S_i^{-1}A) \text{ for all Borel sets } A \subset \mathbb{R}^2.$$

where  $N$  is the total number of contractions.

- Jordan and Rams ('11) computed the **multifractal spectrum** of  $\nu$ ,

$$f_\nu(\alpha) := \dim_{\mathbb{H}}\{x \in \text{supp } \nu : \dim_{\text{loc}}(\nu, x) = \alpha\},$$

building on work of King ('95).

## Theorem (B.–Kolossvary, '21+)

If  $\Lambda$ ,  $\Lambda'$  are Bedford–McMullen carpets with non-uniform vertical fibres, then the intermediate dimensions are equal for all  $\theta$  if and only if the corresponding uniform Bernoulli measures have the same multifractal spectra.

# Bi-Lipschitz equivalence

If  $f: \Lambda \rightarrow \Lambda'$  is bi-Lipschitz then  $\dim_{\theta} \Lambda = \dim_{\theta} \Lambda'$  for all  $\theta$ .

## Corollary

If carpets  $\Lambda$  and  $\Lambda'$  with non-uniform vertical fibres are bi-Lipschitz equivalent then their uniform Bernoulli measures have the same multifractal spectra.

This improves a result of Rao, Yang and Zhang ('21+).

# Thank you for listening!

Please see my poster for more details and to ask any questions!

